

# *Drosophila* Biology in the Genomic Age

Therese Ann Markow<sup>\*,1</sup> and Patrick M. O'Grady<sup>†</sup>

<sup>\*</sup>Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona 85721 and <sup>†</sup>Division of Organisms and the Environment, University of California, Berkeley, California 94720

Manuscript received April 9, 2007

Accepted for publication May 10, 2007

## ABSTRACT

Over the course of the past century, flies in the family Drosophilidae have been important models for understanding genetic, developmental, cellular, ecological, and evolutionary processes. Full genome sequences from a total of 12 species promise to extend this work by facilitating comparative studies of gene expression, of molecules such as proteins, of developmental mechanisms, and of ecological adaptation. Here we review basic biological and ecological information of the species whose genomes have recently been completely sequenced in the context of current research.

If most biologists were given one wish to facilitate their research, many would opt for the fully sequenced genome of their focal taxon. Others might ask for a diverse array of genetic tools around which they could design experiments to answer evolutionary, developmental, behavioral, or ecological questions. With the recent completion of full genome sequences from 12 species, *Drosophila* biologists are now in an unprecedented situation: they have both wishes—and more. Not only does the *Drosophila* model afford researchers full genome sequences and cutting-edge genetic tools, but also more is known about nearly every aspect of the biologies (genetics, development, ecology, phylogenetic relationships, and life history) of these species than of any other eukaryote. Furthermore, because of the comparative genomic framework of the 12 species, discoveries made in one taxon can immediately be placed in a larger evolutionary context.

While most researchers are well aware of the utility of *Drosophila melanogaster* and its close relatives to studies of genetics and developmental biology, few realize that several of the remaining species in this genus have been studied by ecologists and evolutionary biologists nearly since the time that Morgan picked up his first bottle of flies. For example, *D. pseudoobscura*, described by FROLOVA and ASTAUROV (1929), is well known from the classic evolutionary studies of Dobzhansky, his colleagues, and their students (ANDERSON *et al.* 1991; POPADIC and ANDERSON 1994). *D. virilis*, in addition to being a genetic model system in its own right, has also been used to study speciation and chromosome evolution (MCALLISTER 2002; CALETKA and MCALLISTER 2004).

The genus *Drosophila* contains >2000 described species (MARKOW and O'GRADY 2005, 2006), as well as

several hundred taxa that await description. Most of these taxa belong to one of two major subgenera: *Sophophora* and *Drosophila*. Figure 1 shows the phylogenetic relationships and divergence times of the 12 species for which whole-genome sequences are now available. The 12 species with sequenced genomes represent a gradient of evolutionary distances from *D. melanogaster*, including taxa diverging within the past 1 million years to those species who last shared a common ancestor with *D. melanogaster* >30 million years ago (Figure 1). This range was selected to take advantage of the power of multiple, related genomes to discover conserved regulatory motifs, enhance gene prediction, and improve annotation of the *D. melanogaster* genome (BERGMAN *et al.* 2003; BOFFELLI *et al.* 2003). Eight of the newly sequenced species are closely related to *D. melanogaster* and belong to the subgenus *Sophophora*. Five of these, *D. simulans*, *D. yakuba*, *D. erecta*, *D. sechellia*, and *D. ananassae*, are included in the *melanogaster* species group; 2, *D. pseudoobscura* and *D. persimilis*, are placed in the *obscura* group, sister to the *melanogaster* species group; and another, *D. willistoni*, is in the *willistoni* group, a basal clade within *Sophophora* (O'GRADY and KIDWELL 2002). The remaining 3 species belong to the subgenus *Drosophila*, the sister taxon of *Sophophora*. *D. virilis*, a sap flux breeding species, and *D. mojavensis*, a cactophilic taxon, belong to what is referred to as the *virilis-repleta* radiation (THROCKMORTON 1975). *D. grimshawi*, a large, spectacularly patterned species, represents the Hawaiian *Drosophila* radiation, a closely related clade to the *virilis-repleta* species.

Selection of the species to be sequenced thus was based on two criteria: (1) their degree of relatedness to *D. melanogaster* and (2) the likelihood of discovering new genes and new pathways. In the case of the first criterion, it was important to densely sample species closely related to *D. melanogaster* as well as successively more distantly related taxa to discover and annotate conserved regulatory

<sup>1</sup>Corresponding author: Department of Ecology and Evolutionary Biology, BioSciences West 310, 1041 E. Lowell, P.O. Box 210088, Tucson, AZ 85721. E-mail: tmarkow@arizona.edu




FIGURE 1.—(A) Phylogenetic relationships of the 12 fully sequenced *Drosophila* species, along with a timescale for evolution in this group (after Russo *et al.* 1995). Species-level diversity in the containing subgenera and species groups are shown (MARKOW and O'GRADY 2005a and references therein). (B–G) Sex combs in the *melanogaster* species group: (B) *D. melanogaster*, (C) *D. simulans*, (D) *D. sechellia*, (E) *D. erecta*, (F) *D. yakuba*, and (G) *D. ananassae*. (H) Adult male, *D. melanogaster*. (I) Adult male, *D. simulans*. (J) Adult female, *D. ananassae*. (K and L) Sex combs in the *obscura* species group: (K) *D. pseudoobscura* and (L) *D. persimilis*. (M) Adult male, *D. pseudoobscura*. (N) Adult male, *D. virilis*. (O) Adult male, *D. grimshawi*.

regions via phylogenetic shadowing (*e.g.*, BOFFELLI *et al.* 2003). The dense sampling within the *melanogaster* subgroup (*simulans*, *sechellia*, and *yakuba*) and inclusion of the more distantly related *D. erecta* and *D. ananassae* has yielded a much more detailed picture of the *cis*-regulatory regions than comparisons between *melanogaster* and *obscura* (MOSES *et al.* 2006; POLLARD *et al.* 2006). Some of the species were selected because they are behaviorally and ecologically diverse and would yield either novel biochemical pathways or unique variations on already known networks of gene interaction. This avenue has proved particularly relevant for the evolution of the olfactory and gustatory receptor genes in *D. sechellia*, a taxon that oviposits only in the rotting fruit of *Morinda citrifolia*, a highly toxic substrate (MCBRIDE 2007). Another taxon selected on the basis of this criterion is the cactophilic species *D. mojavensis*, in which novel

genes appear to be associated with the use of toxic cactus hosts (MATZKIN *et al.* 2006) as well as with their mating system (KELLEHER and MARKOW 2007).

The first *Drosophila* species, *funnebris*, was described by J. C. Fabricius in 1787 and moved into the genus *Drosophila* by C. F. Fallen in 1823. Meigen described *D. melanogaster* in 1830 (MEIGEN 1830). The number of species described in this group rose slowly throughout the latter half of the 19th century. It was not until the early 1900s, however, after *D. melanogaster* was established as a model organism for understanding genetics that the rate of *Drosophila* species descriptions increased dramatically. Alfred H. Sturtevant, in addition to his contributions to *Drosophila* genetics, also produced early taxonomic treatments of *Drosophila* (STURTEVANT 1916, 1919, 1921, 1939, 1942) and described species such as *D. simulans*, *D. willistoni*, and *D. virilis*. In the late 1930s Th. Dobzhansky

began to use *D. pseudoobscura* and its sibling species, *D. persimilis* and *D. miranda*, in studies aimed at understanding the population genetic basis of species formation. Also around this time, extensive collections by J. T. Patterson and W. S. Stone's group at the University of Texas (Austin, TX) discovered hundreds of new species, mainly from the southwestern United States, Mexico, and Central and South America (PATTERSON 1943; PATTERSON and MAINLAND 1944). Subsequent efforts in Hawaii during the 1960s and the 1970s, the result of a collaboration between the University of Texas group and D. Elmo Hardy at the University of Hawaii at Manoa, discovered an immensely important radiation of *Drosophila* that numbers close to 1000 species (SPIETH 1981). This work led to the popularization of several species as model systems for ecological, population, and behavioral genetics (KAMBYSELLIS 1968; CARSON 1992; DESALLE 1992). Other groups, including those led by Lachaise, Tsacas, David, and Bock, worked throughout the 1970s, 1980s, and 1990s describing many taxa in Africa, Australia, and the South Pacific (BOCK and PARSONS 1981; OKADA 1981; TSACAS *et al.* 1981).

For many problems there is an animal on which it can be most conveniently studied. AUGUST KROGH

According to the Krogh principle, for all biological phenomena there is a perfect model system that can be used to formulate questions and test hypotheses. For example, *D. melanogaster*, popularized by T. H. Morgan and his students in the first half of the 20th century, is one of the premier model systems of modern genetics. Investigations into the ecologies, life histories, and genome features of the 12 fully sequenced species of *Drosophila* demonstrate that each of these flies can be viewed as a model system to address specific biological questions.

In spite of all the work that has been done, what do we really know about the biology and ecology of these 12 species? What makes them compelling models to study interesting questions? What biological questions are each uniquely suited to address? The facility of rearing and manipulating so many diverse but related species of *Drosophila* has fueled the expansion of experimental studies. Several recent reports have reviewed the wealth of research that has been done using *Drosophila* (POWELL 1997; MARKOW and O'GRADY 2005, 2006; ASHBURNER *et al.* 2006). Here, we discuss several areas where the combination of full genome sequences and comparative life history data may help redefine ecological and evolutionary studies.

**Distributions and ecological associations:** Choices about where to feed and oviposit are critical to the survival and fitness of all *Drosophila* species. However, the genetic pathways involved in host plant selection are largely unknown, as are the determinants that make some species specialists and others generalists. In spite of our current lack of understanding about these genes and how they might interact with the environment, they are

of great interest to evolutionary biologists as they may be involved in driving the process of diversification at both the micro- and macroevolutionary levels. Related to host selection and selectivity are those factors (environmental, behavioral, population genetic, and otherwise) that allow some taxa to exist as widespread or cosmopolitan species, while the ranges of others are very narrowly defined. Range maps for *Drosophila* species are given in MARKOW and O'GRADY (2005, 2006). Some species are known to be constrained by host plant distribution and geographic factors, but the ranges, or the bases for the ranges, of others are less well understood.

The 12 species with sequenced genomes display a great diversity in both geographic distribution and ecological association. Some species, such as *D. melanogaster* and *D. simulans*, are cosmopolitan and have spread beyond their ancestral distributions as a result of their commensal association with humans and their ability to breed in a wide variety of rotting fruits. Some close relatives of these generalist species also oviposit in fruit, but are more narrowly distributed and highly selective in their choice of substrate. For example, *D. sechellia* is endemic to the Seychelles and has specialized on the fruits of *M. citrifolia*, a resource toxic to other *Drosophila* (R'KHA *et al.* 1991). Another case of specialization occurs with *D. erecta*, which breeds in species of *Pandanus* in the Ivory Coast of western Africa (LACHAISE and TSACAS 1983). *D. yakuba*, also restricted to Africa, is a generalist fruit breeder (LACHAISE and TSACAS 1983), but has not become a cosmopolitan species like *D. melanogaster* and *D. simulans*. *D. ananassae*, another fruit breeding species that is widespread throughout Asia and the Pacific, is used extensively by some researchers as a genetic model (TOBARI 1992). This species has spread beyond its initial distribution through its association with humans and the fruit trade and is now considered sub-cosmopolitan (SINGH 2000).

The sibling species pair of *D. pseudoobscura* and *D. persimilis* is mainly distributed in western North America, although a small population of *D. pseudoobscura* is located in the mountains near Bogotá, Colombia (DOBZHANSKY *et al.* 1963). During the summer months, both species are abundant in mid- to high-elevation forests, especially those dominated by Ponderosa pines. As temperatures at these sites become colder, populations move to lower elevations and both taxa can be found in or near desert habitats throughout their ranges during the winter. These habitats are not available to these species during the hotter months of the year. Although few breeding records for either species exist, *D. pseudoobscura* has been reared from slime fluxes, domestic fruits, cacti, and agave (POWELL 1997), suggesting that it may be an opportunistic species that can utilize a number of different host types. This would certainly agree with the almost complete lack of overlap in potential host plants between their summer (mountain) and winter (desert) ranges.



*D. willistoni*, a species that breeds in a wide range of rotting fruits, is probably one of the most numerous and broadly distributed drosophilids in the New World and can be found from southern South America to southern North America and throughout the Caribbean (AYALA 1971; DOBZHANSKY and POWELL 1975). Although *D. willistoni* can be readily found in association with humans and the fruit trade within its traditional range, it has not yet been reported outside of the New World.

The three species in the subgenus *Drosophila* that have been sequenced also show a diversity of distributions and ecologies. *D. virilis*, a Holarctic species, has also been reared from fruits in urban settings, but naturally breeds in the fluxes of willows and other decaying parts of trees (THROCKMORTON 1982). *D. mojavensis* is found in the deserts of North America where it breeds in the necroses of several species of cacti (HEED 1978). This species and its relatives have evolved to tolerate not only the toxic compounds found in its hosts, but also the high desiccation conditions of the Sonoran Desert (STRATMAN and MARKOW 1998; GIBBS *et al.* 2003; MATZKIN *et al.* 2006). Although most species of Hawaiian *Drosophila* are highly specific to a single host plant, *D. grimshawi*, a charismatic picture-winged species, is considered a generalist. It utilizes the decaying bark of over seven families of endemic Hawaiian plants (MAGNACCA and O'GRADY 2006).

**Behavioral evolution:** Variability in behavior has been reported for a great many *Drosophila* species, although measures have rarely been made in the same way. Genetics of nonreproductive behaviors for the genus have recently been reviewed by SISODIA and SINGH (2005). Data exist on several of the sequenced species for behaviors such as pupation site preference, locomotor activity, phototaxis, and geotaxis. Far more is known of reproductive behaviors. SPIETH (1952) was the first to categorize the elements of courtship behavior and to describe interspecific variability in these elements. Courtship behaviors of the 12 sequenced species differ in the relative roles of the particular sensory modes in mating: visual, chemical, and auditory (MARKOW and O'GRADY 2005, 2006). For example, *D. melanogaster* and *D. pseudoobscura* will mate equally well in the light and the dark, while mating in their respective sibling species, *D. simulans* and *D. persimilis*, is repressed in darkness. Males of *D. grimshawi*, with their patterned wings, provide elaborate visual displays not seen in males of *D. virilis* or *D. mojavensis*, which tend to focus their courtship activities behind the females. Chemical profiles, or pheromones, differ in close relatives, both within and between species and between the sexes of a given species (FERVEUR 2005). For example, *D. melanogaster* and *D. sechellia* possess long chain dienes not seen in *D. simulans* or *D. erecta*. The longest chain hydrocarbons are seen in *D. mojavensis*. These chemical differences predict that the olfactory and odorant binding receptors of these species would also be different. Finally, species such as *D. virilis* and *D. mojavensis* exhibit both male and female

courtship songs, while in the other 10 species, only the males appear to sing (MARKOW and O'GRADY 2005, 2006; HOIKKALA 2006).

**Life-history evolution:** Ecologists and evolutionary biologists now have the ability to use genomic information and genetic dissection tools to understand the heritable factors contributing to the dazzling array of life-history strategies observed in the genus *Drosophila*. This is an exciting avenue of research that will probe the selective forces that the environment exerts on the genome over evolutionary time. Several developmental and reproductive traits are currently being investigated. For example, there is a clear relationship between body size and egg-to-adult development time: the biggest flies require the longest time to develop (Table 1). Egg-to-adult development time is shortest in *D. melanogaster*, *D. simulans*, and *D. ananassae*, all of which require ~10 days at 24°. The longest development time is in *D. grimshawi*. Development from egg to adult requires nearly a month.

Interspecific differences in reproductive biology represent some of the most interesting features of the 12 species. Few species of *Drosophila* are ready to mate the moment they emerge from the pupa case. In those species in which flies are sexually mature upon emergence, the opposite sex typically requires several days before sexual maturity is achieved. Reproductive maturity times of the 12 species (Table 1) reflect the number of days after emergence by which 80% of flies of a given sex successfully mate with a sexually mature conspecific. Within the subgenus *Sophophora*, adult males tend to mature earlier than females, while in 2 species of the subgenus *Drosophila*, *D. virilis* and *D. mojavensis*, males require two to three time longer to reach sexual maturity. Female *D. grimshawi*, on the other hand, require at least 3 weeks to become sexually mature, almost three times longer than males. Proximate explanations for these differences appear to lie in the relative complexity of either gametogenesis or reproductive tract maturation in one sex or the other. An astounding 15-fold difference in sperm length exists among the 12 species, with *D. persimilis* having the shortest and *D. virilis* the longest sperm. Those species in which males mature before females tend to produce short sperm relative to those in which males mature much later than females (Table 1). In the case of *D. mojavensis*, whose sperm is similar in length to that of *D. melanogaster*, male accessory glands produce relatively large amounts of seminal fluid, the derivatives of which are taken up by females and incorporated into female somatic tissues and developing oocytes (MARKOW and ANKNEY 1984).

The development time in *D. grimshawi* may be related to egg development, rather than sperm formation. *D. grimshawi* females must produce eggs with immensely long chorionic filaments. Kambysellis and his collaborators have shown that the chorionic filament length is adaptive and correlated with the length of the female ovipositor, the type of oviposition substrate, and the depth to

TABLE 1  
Biological characteristics of the 12 sequenced species

Species	F-thorax length (mm)	M-thorax length (mm)	Sexual dimorphism <sup>a</sup>	Egg to adult (days)	F-sexual maturity	M-sexual maturity	Female remating	Ovariole no.	Sperm length
<i>D. melanogaster</i>	0.99 <sup>a</sup>	0.88 <sup>a</sup>	Y <sup>b,c</sup>	10 <sup>a</sup>	4 <sup>a</sup>	2 <sup>a</sup>	5 <sup>a</sup>	43 <sup>a</sup>	1.91 <sup>a</sup>
<i>D. simulans</i>	0.97 <sup>a</sup>	0.87 <sup>a</sup>	Y <sup>b,c</sup>	10 <sup>a</sup>	3 <sup>a</sup>	1 <sup>a</sup>	5 <sup>a</sup>	40 <sup>a</sup>	1.14 <sup>a</sup>
<i>D. sechellia</i>			Y <sup>b,c</sup>	12 <sup>a</sup>				16 <sup>d</sup>	1.60 <sup>e</sup>
<i>D. yakuba</i>			Y <sup>b,c</sup>	11 <sup>a</sup>				28 <sup>e</sup>	1.6 <sup>g</sup>
<i>D. erecta</i>			Y <sup>b,c</sup>	12 <sup>a</sup>				27 <sup>e</sup>	1.2 <sup>g</sup>
<i>D. ananassae</i>	0.94	0.87	Y <sup>b,c</sup>	10 <sup>a</sup>			7	30 <sup>e</sup>	3.3 <sup>g</sup>
<i>D. pseudoobscura</i>	1.09 <sup>a</sup>	1.01 <sup>a</sup>	Y <sup>b,c</sup>	13 <sup>a</sup>	3 <sup>a</sup>	1 <sup>a</sup>	4 <sup>a</sup>	34 <sup>a</sup>	0.36 <sup>a,f</sup>
<i>D. persimilis</i>	1.06 <sup>a</sup>	0.93 <sup>a</sup>	Y <sup>b,c</sup>	13 <sup>a</sup>	4 <sup>a</sup>	0 <sup>a</sup>	4 <sup>a</sup>	36 <sup>a</sup>	0.32 <sup>a,f</sup>
<i>D. willistoni</i>	0.83 <sup>e</sup>	0.79 <sup>e</sup>	N	13 <sup>a</sup>	3 <sup>e</sup>	2 <sup>e</sup>	4 <sup>e</sup>	36 <sup>e</sup>	
<i>D. mojavensis</i>	0.96 <sup>a</sup>	0.89 <sup>a</sup>	N	12 <sup>a</sup>	5 <sup>a</sup>	8 <sup>a</sup>	1 <sup>a</sup>	26 <sup>a</sup>	1.90 <sup>a</sup>
<i>D. virilis</i>	1.33 <sup>a</sup>	1.27 <sup>a</sup>	N	18 <sup>a</sup>	3 <sup>a</sup>	9 <sup>a</sup>	3 <sup>a</sup>	34 <sup>a</sup>	5.70 <sup>a</sup>
<i>D. grimshawi</i>	2.12 <sup>e</sup>	2.23 <sup>e</sup>	N	27 <sup>a</sup>	21 <sup>e</sup>	7 <sup>e</sup>	Rarely <sup>e</sup>	28 <sup>e</sup>	1.19 <sup>e</sup>

Columns 2–9: female (F)-thorax length, male (M)-thorax length, presence of sexual dimorphism, egg-to-adult development time in days, age in days at which 80% of females are sexually receptive, age at which 80% of males are sexually mature, number of days before female remates, female ovariole number, sperm length (mm).

<sup>a</sup> MARKOW and O'GRADY (2005a) and references therein.

<sup>b</sup> Sexual dimorphism for color.

<sup>c</sup> Sexual dimorphism for morphology: male sex combs.

<sup>d</sup> COYNE *et al.* (1991).

<sup>e</sup> T. A. MARKOW (unpublished data).

<sup>f</sup> Sperm length dimorphism, SNOOK (1997).

<sup>g</sup> JOLY and BRESSAC (1974).

which the egg is inserted (KAMBYSELLIS 1993; CRADDOCK and KAMBYSELLIS 1997). As the genetic pathways underlying life-history characteristics are elucidated, biologists will be able to better understand the complex interplay between the genome, development, behavior, and the environment.

**Speciation genetics:** Biologists have long been interested in the genetic changes leading to the formation of new species. Much research has centered on the generation of partial or complete reproductive isolation, both in terms of premating and postmating barriers to the production of viable or fertile offspring. Genetic dissection techniques have been successful in implicating specific chromosomal regions or candidate genes (TING *et al.* 2004; BRIDEAU *et al.* 2006; MOEHRING *et al.* 2006), but the sequenced genomes will allow for finer-scale speciation genetics studies.

Most of the species sequenced have close relatives and have been the subject of intensive studies of speciation genetics involving both interspecific mating experiments and studies using nucleotide variation to examine cases of natural hybridization. When *D. melanogaster* females are crossed with males of either *D. simulans* or *D. sechellia*, the result is the production of sterile females and no males, in accordance with Haldane's rule—meaning that when hybrid inviability or sterility is observed it typically affects the heterogametic sex most profoundly (HALDANE 1922; COYNE 1985; WU *et al.* 1996; ORR 1997). The reciprocal crosses, however, are counter to Haldane's rule in that they produce sterile males and no surviving females (LEMEUNIER *et al.* 1986). Hybrids never have been

obtained between *D. erecta* and any of its relatives, including its closest relative, *D. oreana*, but this is not surprising given the great divergence times between these taxa. In the laboratory, *D. ananassae* produces fertile, viable hybrids in reciprocal crosses with its sibling species *D. pallidosa*, but in nature, sexual isolation, specifically differences in courtship song, prevent the two from interbreeding (YAMADA *et al.* 2002).

Speciation genetics of the *D. pseudoobscura*–*D. persimilis* sibling pair began as early as 1929 (LANCEFIELD 1929). Reciprocal crosses produce sterile male hybrids. With *D. pseudoobscura* mothers, however, ~25% of the F<sub>1</sub> females also are sterile, while the reciprocal cross produces fully fertile females (DOBZHANSKY 1936; ORR 1987). Sequence comparisons of mitochondrial and nuclear genes reveal evidence of recent introgression in different parts of the genomes of these two species (MACHADO and HEY 2003).

The sophophoran most distantly related to *D. melanogaster* is *D. willistoni*. While *D. willistoni* has been reported to inseminate and be inseminated by, at very low levels, its relatives such as *D. equinoxialis* and *D. paulistorum*, its reproductive isolation from these species is effectively complete and no hybrids are produced (BURLA *et al.* 1949). Reproductive barriers also exist within *D. willistoni*. A population of *D. willistoni* collected west of the Andes near Lima, Peru, shows hybrid sterility with *D. willistoni* from the rest of South America, leading AYALA (1972) to designate the Peruvian strains as a separate subspecies, *D. willistoni quecha*.

In the subgenus *Drosophila*, *D. virilis* is able to cross with many of the other species in the *virilis* group. Not

all crosses produce fertile or abundant progeny, however (THROCKMORTON 1982). This suggests that the barriers to reproductive isolation, and therefore the boundaries of what defines a species, in this group may be significantly different from those acting in the subgenus *Sophophora*. The cactophilic sibling pair *D. mojavensis* and *D. arizonae* have become a popular model system for speciation studies because they display a continuum of reproductive isolating mechanisms in interspecific crosses from various populations (MARKOW and HOCUTT 1998). These include premating, postcopulatory–prezygotic, and postzygotic isolation. In addition, the distribution of *D. mojavensis* is bisected by the Sea of Cortez and populations from different regions exhibit signs of incipient speciation (MARKOW and HOCUTT 1998).

Over 95% of the known Hawaiian *Drosophila* species are single island endemics. *D. grimshawi* is unusual in that it occurs on Maui, Molokai, and Lanai. Two of its closest relatives, *D. craddockae* from Oahu and Kauai and *D. pullipes* from Hawaii, are found on the remaining islands and thus the three species are allopatric. Crosses between *D. pullipes* and either *D. craddockae* or *D. grimshawi* produce some viable F<sub>1</sub> progeny, but few males have motile sperm (OHTA 1980), indicating that *D. pullipes* is a distinct species, in spite of only subtle morphological differences. Crosses between *D. grimshawi* and *D. craddockae* produce fertile F<sub>1</sub> progeny, but show a marked reduction in F<sub>2</sub> fertility in reciprocal backcrosses, suggesting evidence for postmating breakdown (KANESHIRO and KAMBYSELLIS 1999). Furthermore, *D. grimshawi* is undergoing differentiation itself, genetically, morphologically, and ecologically, on the different islands it inhabits (PIANO *et al.* 1997).

**The evolution of genome size and rearrangement:** On the basis of comparative cytological studies of metaphase chromosomes, PATTERSON and STONE (1952) suggested that the ancestral karyotype in the genus *Drosophila* is composed of one dot and five acrocentric, or rod, chromosomes (MULLER 1940; STURTEVANT and NOVITSKI 1941; PATTERSON and STONE 1952). All other chromosomal configurations are derived from this basic ancestral state via Robertsonian (ROBERTSON 1957), or centromeric, fusions. It was MULLER (1940) who first hypothesized that the genic content of these six different elements would remain relatively conserved over time because of the rarity of transposition events and the highly deleterious nature of pericentric inversions. The six chromosomal building blocks are lettered A–F and are referred to as Muller's elements. The *D. melanogaster* karyotype of one acrocentric, two metacentric, and one dot chromosome can be generated by two fusion events, one between Muller's B and C and another between Muller's D and E.

Following their role in demonstrating the chromosomal basis of inheritance, *Drosophila* have continued to be a model for studies of genome evolution and rearrangement. Genome sizes (BOSCO *et al.* 2007, accompanying article in this issue; GILBERT 2007) and karyotypes

TABLE 2

Genome sizes from assembled sequences, from flow cytometry (propidium iodide), both in megabases, and chromosome number for each species

Species	Genome size assemblies	Genome size flow cytometry <sup>d</sup>	Chromosome no. ( <i>n</i> )
<i>D. melanogaster</i>	117 <sup>a</sup>	178 ± 16	4
<i>D. simulans</i>	138 <sup>b</sup>	165 ± 2	4
<i>D. sechellia</i>	167 <sup>b</sup>	172 ± 4	4
<i>D. yakuba</i>	166 <sup>b</sup>	187 ± 11	4
<i>D. erecta</i>	153 <sup>b</sup>	146 ± 4	4
<i>D. ananassae</i>	231 <sup>b</sup>	206 ± 9	4
<i>D. pseudoobscura</i>	156 <sup>c</sup>	188 ± 4	4
<i>D. persimilis</i>	188 <sup>b</sup>	188 ± 5	4
<i>D. willistoni</i>	236 <sup>b</sup>	209 ± 7	3
<i>D. mojavensis</i>	193 <sup>b</sup>	143 ± 0	6
<i>D. virilis</i>	206 <sup>b</sup>	313 ± 11	6
<i>D. grimshawi</i>	201 <sup>b</sup>	216 ± 5	6

<sup>a</sup> CELNIKER *et al.* (2002).

<sup>b</sup> GILBERT (2007): <http://insects.eugenes.org/species/>.

<sup>c</sup> RICHARDS *et al.* (2005).

<sup>d</sup> BOSCO *et al.* (2007), accompanying article in this issue.

(Table 2) are quite variable. Gain and loss of heterochromatin are likely explanations for the interspecific differences in genome size (BOSCO *et al.* 2007; GILBERT 2007), while the basic *Drosophila* karyotype of five rods (acrocentric) and a dot chromosome differs among species primarily owing to centromeric fusions.

**Conclusions and prospectus for future research:** The biological diversity of the 12 species provides unparalleled opportunities to address pressing questions about genome evolution, development, behavior, physiology, and species formation. Furthermore, the benefits of the genome sequences are not restricted to the 12 species: for each sequenced taxon, there are multiple related, biologically interesting species for which these genomes will prove a useful and informative springboard to future research. A century after its debut as a research organism, the *Drosophila* model now enters a new era as an even more robust tool for discovery.

We thank Gordon Bennett, Jeremy Bono, Erin Kelleher, Richard Lapoint, Karl Magnacca, Stacy Mazzalupo, Luciano Matzkin, Joel Nitta, Matthew Van Dam, and Tom Watts for critically reading earlier drafts of this manuscript. Kipling Will graciously allowed access to his Microptics camera for the adult and sex comb photos in Figure 1.

#### LITERATURE CITED

- ANDERSON, W. W., J. ARNOLD, D. G. BALDWIN, A. T. BECKENBACH, C. J. BROWN *et al.*, 1991 Four decades of inversion polymorphism in *Drosophila pseudoobscura*. *Proc. Natl. Acad. Sci. USA* **88**: 10367–10371.
- ASHBURNER, M. A., K. G. GOLIC and R. S. HAWLEY, 2006 *Drosophila: A Laboratory Handbook*. Academic Press, New York.
- AYALA, F. J., 1971 Polymorphisms in continental and island populations of *Drosophila willistoni*. *Proc. Natl. Acad. Sci. USA* **68**: 2480–2483.



- AYALA, F. J., 1972 Two new subspecies of the *Drosophila willistoni* group (Diptera). *Pan-Pac. Entomol.* **49**: 273–279.
- BERGMAN, C. M., B. D. PFEIFFER, D. E. RINCON-LIMAS, R. A. HOSKINS, A. GNIRKE *et al.*, 2003 Assessing the impact of comparative genomic sequence data on the functional annotation of the *Drosophila* genome. *Genome Biol.* **3**: 1–20.
- BOCK, I. R., and P. A. PARSONS, 1981 Species of Australia and New Zealand, pp. 291–308 in *The Genetics and Biology of Drosophila*, Vol. 3a, edited by M. A. ASHBURNER, H. L. CARSON and J. N. THOMPSON. Academic Press, London.
- BOFFELLI, D., J. McAULIFFE, D. OVCHARENKO, K. D. LEWIS, I. OVCHARENKO *et al.*, 2003 Phylogenetic shadowing of primate sequences to find functional regions of the human genome. *Science* **299**: 1391–1394.
- BOSCO, G., P. CAMPBELL, J. T. LEIVA-NETO and T. A. MARKOW, 2007 Analysis of *Drosophila* species genome size and satellite DNA content reveals differences among strains as well as between species. *Genetics* **177**: 1277–1290.
- BRIDEAU, N. J., H. A. FLORES, J. WANG, S. MAHESHWARI, X. WANG *et al.*, 2006 Two Dobzhansky-Muller genes interact to cause hybrid lethality in *Drosophila*. *Science* **314**: 1292–1295.
- BURLA, H., A. BRITO DA CUNHA, A. R. CORDERO, TH. DOBZHANSKY, C. MALAGOLOWKIN *et al.*, 1949 The *willistoni* group of sibling species of *Drosophila*. *Evolution* **3**: 300–314.
- CALETKA, B. C., and B. F. McALLISTER, 2004 A genealogical view of chromosomal evolution and species delimitation in the *Drosophila virilis* species subgroup. *Mol. Phylogenet. Evol.* **33**: 664–670.
- CARSON, H. L., 1992 Inversions in Hawaiian *Drosophila*, pp. 407–439 in *Drosophila Inversion Polymorphism*, edited by C. B. KRIMBAS and J. R. POWELL. CRC Press, Boca Raton, FL.
- COYNE, J. A., 1985 The genetic basis of Haldane's rule. *Nature* **314**: 736–738.
- COYNE, J. A., J. RUX and J. R. DAVID, 1991 Genetics of morphological differences and hybrid sterility between *Drosophila sechellia* and its relatives. *Genet. Res.* **57**: 113–122.
- CELNIKER, S. E., E. FRISE, A. HODGSON, R. A. GEORGE, R. A. HOSKINS *et al.*, 2002 Finishing a whole-genome shotgun: release 3 of the *Drosophila melanogaster* euchromatic genome sequence. *Genome Biol.* **3**: 1–14.
- CRADDOCK, E. M., and M. P. KAMBYSELLIS, 1997 Adaptive radiation in the Hawaiian *Drosophila* (Diptera: Drosophilidae): ecological and reproductive character analyses. *Pac. Sci.* **51**(4): 475–489.
- DESALLE, R., 1992 The intra- and inter-island relationships of Hawaiian *Drosophila* deduced from DNA sequence information. *Am. J. Bot.* **79**(Suppl. 6): 126.
- DOBZHANSKY, TH., 1936 Studies of hybrid sterility II. Localization of sterility factors in *D. pseudoobscura* hybrids. *Genetics* **21**: 113–135.
- DOBZHANSKY, TH., and J. R. POWELL, 1975 The willistoni group of sibling species of *Drosophila*, pp. 589–622 in *Invertebrates of Genetic Interest*, edited by R. C. KING. Plenum Press, New York.
- DOBZHANSKY, TH., A. S. HUNTER, O. PAVLOVSKY, B. SPASSKY and B. WALLACE, 1963 Genetics of natural populations. XXXI. Genetics of an isolated marginal population of *Drosophila pseudoobscura*. *Genetics* **48**: 91–103.
- FERVEUR, J. F., 2005 Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal communication. *Behav. Genet.* **35**: 279–295.
- FROLOVA, S. L., and B. L. ASTUAROV, 1929 Die chromosomengarnitur als systematisches Merkmal. Eine verleichende untersuchung der russischen und amerikansichen *Drosophila obscura* Fallen. *Z. Zellforsch. Mikrosk. Anat.* **10**: 201–213.
- GIBBS, A. G., M. C. PERKINS and T. A. MARKOW, 2003 No place to hide: microclimates of desert *Drosophila*. *J. Therm. Biol.* **28**: 353–362.
- GILBERT, D. G., 2007 DroSpeGe: rapid access database for new *Drosophila* species genomes. *Nucleic Acids Res.* **35**: D480–D485.
- HALDANE, J. B. S., 1922 Sex-ratio and unisexual sterility in hybrid animals. *Genetics* **12**: 101–109.
- HEED, W. B., 1978 Ecology and genetics of Sonoran Desert *Drosophila*, pp. 109–126 in *Symposium on Genetics and Ecology: The Interface*, edited by P. F. BRUSSARD. Springer-Verlag, Heidelberg, Germany.
- HOIKKALA, A., 2005 Inheritance of male sound characteristics in *Drosophila* species, in *Insect Sounds and Communication; Physiology, Ethology and Evolution*, edited by S. DROSPOULIS and M. CLARIDGE. Taylor & Francis, Boca Raton, FL.
- JOLY D., C. and BRESSAC, 1994 Sperm length in Drosophilidae (Diptera): estimation by testis and receptacle lengths. *Int. J. Insect Morph. Embryol.* **23**(2): 85–92.
- KAMBYSELLIS, M. P., 1968 Comparative studies of oogenesis and egg morphology among species of the genus *Drosophila*. *Univ. Tex. Publ. Stud. Genet.* **4**(6818): 71–92.
- KAMBYSELLIS, M. P., 1993 Ultrastructural diversity in the egg chorion of Hawaiian *Drosophila* and *Scaptomysca*: ecological and phylogenetic considerations. *Int. J. Insect Morph. Embryol.* **22**: 417–446.
- KANESHIRO, K. Y., and M. P. KAMBYSELLIS, 1999 Description of a new allopatric sibling species of Hawaiian picture-winged *Drosophila*. *Pac. Sci.* **53**(2): 208–213.
- KELLEHER, E. S., and T. A. MARKOW, 2007 Reproductive tract interactions contribute to isolation in *Drosophila*. *Fly* **1**: 33–37.
- LANCEFIELD, D. E., 1929 A genetic study of crosses of two races or physiological species of *Drosophila obscura*. *Z. Indukt. Abstammungs. Vererbungs.* **52**: 287–317.
- LEMEUNIER, F., J. R. DAVID, L. TSACAS and M. A. ASHBURNER, 1986 The *melanogaster* species group, pp. 147–256 in *The Genetics and Biology of Drosophila*, Vol. 3e, edited by M. A. ASHBURNER, H. L. CARSON and J. N. THOMPSON. Academic Press, London.
- MACHADO, C. A., and J. HEY, 2003 The causes of phylogenetic conflict in a classic *Drosophila* species group. *Proc. R. Soc. Lond. Ser. B* **270**: 1193–1202.
- MAGNACCA, K. N., and P. M. O'GRADY, 2006 A subgroup structure for the modified mouthparts group of Hawaiian *Drosophila*. *Proc. Hawaii. Entomol. Soc.* **38**: 87–101.
- MARKOW, T. A., and P. F. ANKNEY, 1984 *Drosophila* males contribute to oogenesis in a multiple mating species. *Science* **224**: 302–303.
- MARKOW, T. A., and G. D. HOCUTT, 1998 Speciation in Sonoran Desert *Drosophila*: testing the limits of the rules, pp. 234–244 in *Endless Forms: Species and Speciation*, edited by D. A. HOWARD and S. BERLOCHER. Oxford University Press, New York.
- MARKOW, T. A. and P. M. O'GRADY, 2005 *Drosophila: A Guide to Species Identification and Use*. Academic Press, London.
- MARKOW, T. A., and P. M. O'GRADY, 2006 Evolutionary genetics of reproductive behavior in *Drosophila*. *Annu. Rev. Genet.* **39**: 263–291.
- MATZKIN, L. M., T. WATTS, B. G. BITLER, C. A. MACHADO and T. A. MARKOW, 2006 Functional genomics of cactus host shifts in *Drosophila mojavensis*. *Mol. Ecol.* **15**: 4635–4643.
- McALLISTER, B. F., 2002 Chromosomal and allelic variation in *Drosophila americana*: selective maintenance of a chromosomal cline. *Genome* **45**: 13–21.
- MCBRIDE, C. S., 2007 Rapid evolution of smell and taste receptor genes during host specialization in *Drosophila sechellia*. *Proc. Natl. Acad. Sci. USA* **104**: 4996–5001.
- MEIGEN, J. W., 1830 *Systematische Beschreibung der Bekannten Europäischen Zweiflügeligen Insekten*. Theil. Schulz, Hamm, Germany.
- MOEHRING, A. J., A. LLOPART, S. ELWYN, J. A. COYNE and T. F. MACKAY, 2006 The genetic basis of postzygotic reproductive isolation between *Drosophila santomea* and *D. yakuba*. *Genetics* **173**: 225–233.
- MOSES, A. M., D. A. POLLARD, D. A. NIX, V. N. IYER, X. LI *et al.*, 2006 Large-scale turnover of functional transcription factor binding sites in *Drosophila*. *PLoS Comput. Biol.* **2**: 1219–1231.
- MULLER, H. J., 1940 Bearings of the *Drosophila* work on systematics, pp. 185–268 in *The New Systematics*, edited by J. HUXLEY. Clarendon Press, Oxford.
- O'GRADY, P. M., and M. G. KIDWELL, 2002 Phylogeny of the subgenus *sophophora* (Diptera: Drosophilidae) based on combined analysis of nuclear and mitochondrial sequences. *Mol. Phylogenet. Evol.* **22**(3): 442–453.
- OHTA, A., 1980 Coadapted gene complexes in incipient species of Hawaiian *Drosophila*. *Am. Nat.* **115**: 121–132.
- OKADA, T., 1981 Oriental species, including New Guinea, pp. 261–290 in *The Genetics and Biology of Drosophila*, Vol. 3a, edited by M. A. ASHBURNER, H. L. CARSON and J. N. THOMPSON. Academic Press, London.
- ORR, H. A., 1987 Genetics of male and female sterility in hybrids of *Drosophila pseudoobscura* and *D. persimilis*. *Genetics* **116**: 555–565.
- ORR, H. A., 1997 Haldane's rule. *Annu. Rev. Evol. Syst.* **28**: 195–218.
- PATTERSON, J. T., 1943 The *Drosophila* of the Southwest. *Univ. Tex. Publ.* **4313**: 7–216.
- PATTERSON, J. T., and G. B. MAINLAND, 1944 The Drosophilidae of Mexico. *Univ. Tex. Publ.* **4445**: 9–101.

- PATTERSON, J. T., and W. S. STONE, 1952 *Evolution in the Genus Drosophila*. Macmillian Press, New York.
- PIANO, F., E. M. CRADDOCK and M. P. KAMBYSELLIS, 1997 Phylogeny of the island populations of the Hawaiian *Drosophila grimshawi* complex: evidence of combined data. *Mol. Phylogenet. Evol.* **7**: 173–184.
- POLLARD, D. A., A. M. MOSES, V. N. IYER and M. B. EISEN, 2006 Detecting the limits of regulatory element conservation and divergence estimation using pairwise and multiple alignments. *BMC Bioinform.* **7**: 376–390.
- POPADIC, A., and W. W. ANDERSON, 1994 The history of a genetic system. *Proc. Natl. Acad. Sci. USA* **91**: 6819–6823.
- POWELL, J. R., 1997 *Progress and Prospects in Evolutionary Biology: The Drosophila Model*. Oxford University Press, New York.
- RICHARDS, S., R. P. MEISEL, O. COURONNE, S. HUA, M. A. SMITH *et al.*, 2005 Comparative genome sequencing of *Drosophila pseudoobscura*: chromosomal, gene, and cis-element evolution. *Genome Res.* **15**: 1–18.
- R'KHA, S., P. CAPY and J. R. DAVID, 1991 Host plant specialization in the *Drosophila melanogaster* species complex: a physiological, behavioral and genetic analysis. *Proc. Natl. Acad. Sci. USA* **88**: 1835–1839.
- ROBERTSON, F. W., 1957 Studies in quantitative inheritance. *J. Genet.* **55**: 410–427.
- RUSO, C. A. M., N. TAKEZAKI and M. NEI, 1995 Molecular phylogeny and divergence times of drosophilid species. *Mol. Biol. Evol.* **12**(3): 391–404.
- SINGH, B. N., 2000 *Drosophila ananassae*: a species characterized by several unusual genetic features. *Curr. Sci.* **78**: 391–398.
- SISODIA, S., and B. N. SINGH, 2005 Behaviour genetics of *Drosophila*: non-sexual behaviour. *J. Genet.* **84**: 195–216.
- SNOOK, R. R., 1997 Is the production of multiple sperm types adaptive? *Evolution* **51**(3): 797–808.
- SPIETH, H. T., 1952 Mating behaviour within the genus *Drosophila* (Diptera). *Bull. Am. Mus. Nat. Hist.* **99**: 395–474.
- SPIETH, H. T., 1981 History of the Hawaiian *Drosophila* project. *DIS* **56**: 6–14.
- STRATMAN, R., and T. A. MARKOW, 1998 Resistance to thermal stress in desert *Drosophila*. *Funct. Ecol.* **8**: 965–970.
- STURTEVANT, A. H., 1916 Notes on North American Drosophilidae with descriptions of 23 new species. *Ann. Entomol. Soc. Am.* **9**: 323–343.
- STURTEVANT, A. H., 1919 A new species closely resembling *Drosophila melanogaster*. *Psyche* **26**: 153–155.
- STURTEVANT, A. H., 1921 The North American species of *Drosophila*. *Publ. Carnegie Inst.* **301**: 1–150.
- STURTEVANT, A. H., 1939 On the subdivision of the genus *Drosophila*. *Proc. Natl. Acad. Sci. USA* **25**: 137–141.
- STURTEVANT, A. H., 1942 The classification of the genus *Drosophila*, with descriptions of nine new species. *Univ. Tex. Publ.* **4213**: 5–51.
- STURTEVANT, A. H., and E. NOVITSKI, 1941 The homologies of the chromosome elements in the genus *Drosophila*. *Genetics* **26**: 517–541.
- THROCKMORTON, L. H., 1975 The phylogeny, ecology and geography of *Drosophila*, pp. 421–469 in *Invertebrates of Genetic Interest* (Handbook of Genetics, Vol. 3), edited by R. C. KING. Plenum Press, New York.
- THROCKMORTON, L. H., 1982 The *virilis* species group, pp. 227–297 in *The Genetics and Biology of Drosophila*, Vol. 3b, edited by M. A. ASHBURNER, H. L. CARSON and J. N. THOMPSON. Academic Press, London.
- TING, C. T., S. C. TSAUR, S. SUN, W. E. BROWNE, Y. C. CHEN *et al.*, 2004 Gene duplication and speciation in *Drosophila*: evidence from the Odysseus locus. *Proc. Natl. Acad. Sci. USA* **101**: 12232–12235.
- TOBARI, Y. N., 1992 *Drosophila ananassae*. *Genetical and Biological Aspects*. Japan Scientific Societies Press, Tokyo.
- TSACAS, L., D. LACHAISE and J. R. DAVID, 1981 Composition and biogeography of the Afrotropical Drosophilid fauna, pp. 197–260 in *The Genetics and Biology of Drosophila*, Vol. 3a, edited by M. A. ASHBURNER, H. L. CARSON and J. N. THOMPSON. Academic Press, London.
- WU, C. I., N. A. JOHNSON and M. F. PALOPOLI, 1996 Haldane's rule and its legacy: Why are there so many sterile males? *Trends Ecol. Evol.* **11**: 281–284.
- YAMADA, H., M. MATSUDA and Y. OGUMA, 2002 Genetics of sexual isolation based on courtship song between two sympatric species: *Drosophila ananassae* and *D. pallidosa*. *Genetica* **116**: 225–237.

Communicating editor: R. S. HAWLEY